

477 W. Horton Rd. Bellingham, WA 98226 Phone (360) 933-4892 Fax (360) 933-1197

Patient Name:	
Date of Birth:	Weight:
IV Access:	Height:
Allergies:	

## **Iron Order Form**

Please complete this form and fax to (360)933-1197. Call our office and our clinical pharmacists will be happy to make therapy recommendations.

<u>Diagnoses:</u>		☐ Iron Deficiency Anemia second Iron Deficiency Anemia second Unspecified Iron Deficiency Anemia second Iron Deficiency Anemia Iron Deficiency Anemia Iron Deficiency Anemia Iron Deficiency Anemia Iron Iron Deficiency Anemia Iron Iron Iron Iron Iron Iron Iron Iron	ondary to inadequate dietary intake	ICD-10: D50.0 ICD-10: D50.8 ICD-10: D50.9		
		☐ Other:		ICD-10:		
Scree	Screening: Does patient have a history of: □drug allergies □asthma □autoimmune disorder					
	Is the patient pregnant?    Yes   No					
Medic	ation C					
	Iron Su	crose (Venofer): mg IV	every days for	doses.		
	(Recommend 100-300 mg per dose, and 1,000 mg per course; optimal frequency is ≤ 3 times weekly)					
	Ferric Carboxymaltose (Injectafer): 🗖 15 mg/kg (max 750 mg) IV every 7 days for 2 doses					
	□ Alternate instructions:					
	Other formulation:					
* *						
Nursing Orders:						
•	Obtain vital signs before start of therapy.					
•	Observe for hypotension and have Infusion Reaction Management kit with NS immediately available.					
•	RN to insert Peripheral IV, rotate sites as needed, and remove after completion of therapy.					
	Other:					
<u>Labs:</u>	•					
	CBC w		☐ 1 week after therapy completion			
	Serum TIBC (i	ncludes iron & transferritin sat.)	<ul><li>☐ 1 week after therapy completion</li><li>☐ 1 week after therapy completion</li></ul>			
	`	,	every			
_						
P	rescriber S	ignature	Date			
_						
P	lease Print	Name				

Page 1 of 1

**KEY:** • Orders are initiated unless crossed out by provider.