

477 W. Horton Rd. Bellingham, WA 98226 Phone (360) 933-4892 Fax (360) 933-1197

Patient Name:	
Date of Birth:	Weight:
IV Access:	Height:
Allergies:	

Stelara (for Crohn's or UC) Order Form

Please fax this form, copies of insurance cards, demographics, and supporting clinical documentation to (360) 933-1197 to facilitate an efficient referral. Thank you for choosing Infusion Solutions!

<u>Diagnos</u>	es: Crohn's Disease Ulcerative Colitis Other:	ICD-10: ICD-10: ICD-10:		
TB Histo	Dry: Date of last PPD test:	Result:		
Medication Orders:				
♦ S ²	telara (ustekinumab)			
	 Induction: Administer IV over at least one hour as a single dose. Use 0.2 micron in-line filter. Dose: □260 mg (<= 55 kg) □390 mg (>55 to 85 kg) □520 mg (>85 kg) Maintenance: Inject 90 mg subcutaneously every 8 weeks (starting 8 weeks after IV induction dose). 			
♦ Fl ♦ Li	 Alteplase 2mg IV to declot central IV access per Infusion Solutions protocol as needed for occlusion. Flush line with D5W, 0.9% NaCl and/or Heparin 10 units/ml or 100 units/ml per Infusion Solutions protocol. Lidocaine 1% - up to 0.2ml intradermally PRN (may buffer with sodium bicarbonate 8.4% in 10:1 ratio). Infusion Reaction Management per Infusion Solutions protocol as needed. 			
Nursing	Orders:			
◆ M ◆ R	no central IV access, RN to insert peripheral IV, rotate s lonitor for infusion reactions during IV induction dose, an N to administer subQ injections, and train patient to self other:	nd for 30 minutes after infusion. f-inject if deemed clinically appropriate.		
Labs:				
□ c	BC with differential every			
		every		
		every		

Prescriber Signature

Date

Please Print Name

KEY: • Orders are initiated unless crossed out by provider.

Check box to initiate order.