

477 W. Horton Rd. Bellingham, WA 98226 Phone (360) 933-4892 Fax (360) 933-1197

Patient Name:	
Date of Birth:	Weight:
IV Access:	Height:
Allergies:	

Entyvio (Vedolizumab) Order Form

Please fax this form, copies of insurance cards, demographics, and supporting clinical documentation to (360) 933-1197 to facilitate an efficient referral. Thank you for choosing Infusion Solutions!

Diagnose	es:	IC	D-10:	
	Ulcerative colitis	IC	D-10:	
	☐ Other:	IC	D-10:	
	nt received Entyvio before? nunizations up to date?	□ No□ Yes (date of last infusion□ Yes□ No (recommended before)	n:) re initiating therapy – please indicate	
Medication	on Orders:			
• E	ntyvio (Vedolizumab)			
	☐ 300 mg IV over 30 minutes at 0, 2, and 6 weeks, then every 8 weeks thereafter			
	☐ Other dose/instructions:			
 Alteplase 2mg IV to declot central IV access per Infusion Solutions protocol as needed for occlusion. Flush line with 0.9% NaCl, D5W and/or Heparin 10 units/ml or 100 units/ml per Infusion Solutions protocol. Lidocaine 1% - up to 0.2ml intradermally PRN (may buffer with sodium bicarbonate 8.4% in 10:1 ratio). Infusion Reaction Management per Infusion Solutions protocol as needed. 				
Nursing Orders:				
 If no central IV access, RN to insert peripheral IV, rotate site as needed, and remove at end of therapy. Monitor for infusion/hypersensitivity reactions during infusions. Monitor for signs of infection or liver impairment before each infusion; contact prescriber if infection is present (dose may be held). Other: 				
<u>Labs:</u> □	CBC w/diff	□Each infusion	☐Other frequency	
	CMP	□Each infusion	Other frequency	
	Hepatic function panel	□Each infusion □Each infusion	☐Other frequency	
	Other:	□Each infusion	□Other frequency	
Presci	iber Signature		Date	
Please	Print Name			

KEY: ♦ Orders are initiated unless crossed out by provider.

☐ Check box to initiate order.